```
Order Coleoptera Linnaeus, 1758 (4 suborders)<sup>1, 2</sup>
```

†Suborder **Protocoleoptera** Tillyard, 1934 (3 superfamilies)

†Superfamily **Tshekardocoleoidea** Rohdendorf, 1944 (3 families)

†Family **Tshekardocoleidae** Rohdendorf, 1944 (13 genera, 18 species)

†Family Labradorocoleidae Ponomarenko, 1969 (1 genus, 1 species)

†Family **Oborocoleidae** Kukalová, 1969 (2 genera, 2 species)

†Superfamily **Permocupedoidea** Martynov, 1933 (2 families)

†Family **Permocupedidae** Martynov, 1933 (11 genera, 30 species)

†Family **Taldycupedidae** Rohdendorf, 1961 (6 genera, 18 species)

†Superfamily **Permosynoidea** Tillyard, 1924 (2 families)

†Family Ademosynidae Ponomarenko, 1968 (12 genera, 29 species)

†Family **Permosynidae** Tillyard, 1924 (3 genera, 14 species)

## Suborder Archostemata Kolbe, 1908 (8 families)

Family Crowsoniellidae Iablokoff-Khnzorian, 1983 (1 genus, 1 species)

Family Cupedidae Laporte, 1836 (53(†44) genera, 120(†89) species)

Family **Micromalthidae** Barber, 1913 (2(†1) genera, 2(†1) species)

Family **Ommatidae** Sharp and Muir, 1912 (10(†8) genera, 120(†114) species)

Family **Jurodidae** Ponomarenko, 1985 (2(†1) genera, 3(†2) species)

†Family **Triadocupedidae** Ponomarenko, 1966 (8 genera, 16 species)

†Family Magnocoleidae Hong, 1998 (1 genus, 1 species)

†Family **Obrieniidae** Zherikhin and Gratshev, 1994 (5 genera, 7 species)

# Suborder Myxophaga Crowson, 1955 (5 superfamilies)

†Superfamily Asiocoleoidea Rohdendorf, 1961 (2 families)

†Family Asiocoleidae Rohdendorf, 1961 (1 genus, 1 species)

†Family **Tricoleidae** Ponomarenko, 1969 (3 genera, 6 species)

†Superfamily Rhombocoleoidea Rohdendorf, 1961 (1 family)

†Family Rhombocoleidae Rohdendorf, 1961 (5 genera, 10 species)

Superfamily Schizophoroidea Ponomarenko, 1968 (3 families)

†Family Schizophoridae Ponomarenko, 1968 (20 genera, 36 species)

†Family Catiniidae Ponomarenko, 1968 (5 genera, 6 species)

†Family Schizocoleidae Rohdendorf, 1961 (4 genera, 30 species)

Superfamily Lepiceroidea Hinton, 1936 (1882) (2 families)

Family Lepiceridae Hinton, 1936 (1882) (1(†1) genus, 2(†1) species)<sup>3</sup>

Superfamily Sphaeriusoidea Erichson, 1845 (3 families)

Family **Torridincolidae** Steffan, 1964 (7 genera, 60 species)

Family Hydroscaphidae LeConte, 1874 (3 genera, 22 species)<sup>4</sup>

Family **Sphaeriusidae** Erichson, 1845 (2 (†1) genus, 20 (†1) species)

## Suborder Adephaga Schellenberg, 1806 (17 families)

†Family Tritarsusidae Hong, 2002 (1 genus, 1 species)

Family Gyrinidae Latreille, 1810 (25 (†13) genera, 1000 (†118) species)

Family Trachypachidae C. G. Thomson, 1857 (22 (†21) genera, 57 (†51) species)

Family Meruidae Spangler and Steiner, 2005 (1 genus, 1 species)

Family **Rhysodidae** Laporte, 1840 (40 genera, 350 species)

Family Carabidae Latreille, 1802 (1500 genera, 40000 species)

See Fikáček and Šípková (2009).

Zootaxa 3148 © 2011 Magnolia Press · 203

<sup>1.</sup> **BY** S. A. Slipinski, R. A. B. Leschen, and J. F. Lawrence (for full contact details, see the list after **References**). The title of this contribution should be cited as "Order Coleoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness".

<sup>2.</sup> Recent Coleoptera currently includes 176 families, 29500 genera and 386500 species, fossil taxa includes 31 families, 230 genera and 600 species. The classification is based on Lawrence and Newton (1995) and Bouchard *et al.* (2011). Numbers for fossil taxa were derived from Kirejtshuk et al. (2010) and only numbers of fossils for extinct Protocoleoptera and Coleoptera families are provided, apart from extant primitive Polyphaga. Numbers for extant taxa were taken from the *Handbook of Zoology* chapters in Beutel & Leschen (2005) and Leschen et al. (2010) except where noted. Because up-to-date catalogues do not exist, numbers in the *Handbook* had included estimates and here we cross-referenced these estimates with recent catalogues (e.g., Löbl & Smetana 2007) and web-based information (e.g., Hallan 2010). Cut off date for the information contained here was 10 February 2011.

<sup>3.</sup> See Ge et al. (2010).

```
Family Haliplidae Aubé, 1836 (5 (†1) genera, 220 (†2) species)
               †Family Triaplidae Ponomarenko, 1977 (1 genus, 2 species)
               †Family Colymbotethidae Ponomarenko, 1994 (2 genera, 2 species)
               †Family Parahygrobiidae Ponomarenko, 1977 (1 genus, 1 species)
               †Family Coptoclavidae Ponomarenko, 1961 (14 genera, 32 species)
               †Family Liadytidae Ponomarenko, 1977 (2 genera, 6 species)
               Family Noteridae C. G. Thomson, 1859 (14 genera, 250 species)
               Family Amphizoidae LeConte, 1853 (1 genus, 5 species)
               Family Aspidytidae Ribera et al., 2002 (1 genus, 2 species)
               Family Hygrobiidae Régimbart, 1879 (1837) (1 genus, 5 species)
               Family Dytiscidae Leach, 1815 (190 (†6) genera, 4100 (†85) species)
Suborder Polyphaga Emery, 1886 (7 series)
     Series Staphyliniformia Latreille, 1802 (2 superfamilies)
          Superfamily Hydrophiloidea Latreille, 1802 (4 families)
               Family Hydrophilidae Latreille, 1802 (200 genera, 3400 species)
               Family Sphaeritidae Shuckard, 1839 (1 genus, 5 species)
               Family Synteliidae Lewis, 1882 (1 genus, 7 species)
               Family Histeridae Gyllenhal, 1808 (350 genera, 4300 species)
          Superfamily Staphylinoidea Latreille, 1802 (6 families)
               Family Hydraenidae Mulsant, 1844 (42 genera, 1600 species)
               Family Ptiliidae Erichson, 1845 (80 genera, 650 species)
               Family Agyrtidae C. G. Thomson, 1859 (8 genera, 70 species)
               Family Leiodidae Fleming, 1821 (342 genera, 3700 species)
               Family Silphidae Latreille, 1807 (15 genera, 200 species)
               Family Staphylinidae Latreille, 1802 (3500 genera, 56000 species)
    Series Scarabaeiformia Crowson 1960 (1 superfamily)
          Superfamily Scarabaeoidea Latreille, 1802 (14 families)
               Family Pleocomidae LeConte, 1861 (2 genera, 50 species)
               Family Geotrupidae Latreille, 1802 (70 genera, 920 species)
               Family Belohinidae Paulian, 1959 (1 genus, 1 species)
              Family Passalidae Leach, 1815 (64 genera, 800 species)
               Family Trogidae W. S. MacLeay, 1819 (5 genera, 300 species)
              Family Glaresidae Kolbe, 1905 (3(†2) genera, 60 (†3) species)
               Family Diphyllostomatidae Holloway, 1972 (1 genus, 3 species)
               Family Lucanidae Latreille, 1804 (120(†6) genera, 1500 (†11) species)
               Family Ochodaeidae Mulsant and Rey, 1871 (15 genera, 110 species, †2/3)
               Family Hybosoridae Erichson, 1847 (77(†3) genera, 580 (†7) species)
               Family Glaphyridae W. S. MacLeay, 1819 (11(†1) genera, 210(†6) species)
               Family Scarabaeidae Latreille, 1802 (1900 genera, 27000 species)
               †Family Coprinisphaeridae Genise, 2004 (4 genera, 9 species, ichnotaxon)
               †Family Pallichnidae Genise, 2004 (2 genera, 3 species, ichnotaxon)
    Series Scirtiformia Fleming, 1821 (1 superfamily)
          Superfamily Scirtoidea Fleming, 1821 (6 families)
               Family Scirtidae Fleming, 1821 (35 genera, 800 species)
               Family Decliniidae Nikitsky et al., 1994 (1 genus, 2 species)
               Family Eucinetidae Lacordaire, 1857 (11 genera, 53 species)
               Family Clambidae Fischer von Waldheim, 1821 (6 genera, 170 species)
               †Family Elodophthalmidae Kirejtshuk and Azar, 2008 (1 genus, 2 species)
               †Family Mesocinetidae Kirejtshuk and Ponomarenko, 2010 (4 genera, 7 species)
    Series Elateriformia Crowson 1960 (4 superfamilies)
          Superfamily Dascilloidea Guérin-Méneville, 1843 (1834) (2 families)
               Family Dascillidae Guérin-Méneville, 1843 (1834) (15 genera, 80 species)
               Family Rhipiceridae Latreille, 1834 (7 genera, 70 species)
          Superfamily Buprestoidea Leach, 1815 (1 family)
```

```
Family Buprestidae Leach, 1815 (470 genera, 14700 species)
     Superfamily Byrrhoidea Latreille, 1804 (12 families)
          Family Byrrhidae Latreille, 1804 (38 genera, 430 species)
          Family Elmidae Curtis, 1830 (149 genera, 1500 species)
          Family Dryopidae Billberg, 1820 (1817) (33 genera, 300 species)
          Family Lutrochidae Kasap and Crowson, 1975 (1 genus, 11 species)
          Family Limnichidae Erichson, 1846 (37 genera, 390 species)
          Family Heteroceridae W. S. MacLeay, 1825 (15 genera, 300 species)
          Family Psephenidae Lacordaire, 1854 (35 genera, 290 species)
          Family Cneoglossidae Champion, 1897 (1 genus, 10 species)
          Family Ptilodactylidae Laporte, 1836 (34 genera, 500 species)
          Family Chelonariidae Blanchard, 1845 (3 genera, 250 species)
          Family Eulichadidae Crowson, 1973 (2 genera, 30 species)
          Family Callirhipidae Emden, 1924 (9 genera, 150 species)
     Superfamily Elateroidea Leach, 1815 (20 families)
          Family Rhinorhipidae Lawrence, 1988 (1 genus, 1 species)
          Family Artematopodidae Lacordaire, 1857 (8 genera, 45 species)
          Family Brachypsectridae LeConte and G. H. Horn, 1883 (1 genus, 5 species)
          Family Cerophytidae Latreille, 1834 (3 genera, 21 species)
          Family Eucnemidae Eschscholtz, 1829 (200 genera, 1500 species)
          Family Throscidae Laporte, 1840 (5 genera, 150 species)
          Family Elateridae Leach, 1815 (400 genera, 10000 species)
          Family Plastoceridae Crowson, 1972 (1 genus, 2 species)
          Family Drilidae Blanchard, 1845 (6 genera, 120 species)
          Family Omalisidae Lacordaire, 1857 (3 genera, 8 species)
          Family Lycidae Laporte, 1836 (160 genera, 4600 species)
          Family Telegeusidae Leng, 1920 (3 genera, 10 species)
          Family Phengodidae LeConte, 1861 (31 genera, 250 species)
          Family Rhagophthalmidae E. Olivier, 1907 (6 genera, 30 species)<sup>5</sup>
          Family Lampyridae Rafinesque, 1815 (110 genera, 2200 species)
          Family Omethidae LeConte, 1861 (8 genera, 33 species)
          Family Cantharidae Imhoff, 1856 (1815) (160 genera, 5100 species)
          Family Podabrocephalidae Pic. 1930 (1 genus. 1 species)
          †Family Berendtimiridae Winkler, 1987 (1 genus, 1 species)
          †Family Lasiosynidae Kirejtshuk, Chang, Ren & Kun, 2010 (4 genera, 7 species)
Series Derodontiformia LeConte, 1861 (1 superfamily)
     Superfamily Derodontoidea LeConte, 1861 (3 families)
          Family Derodontidae LeConte, 1861 (4 genera, 30 species)
          Family Nosodendridae Erichson, 1846 (1 genus, 50 species)
          Family Jacobsoniidae Heller, 1926 (3 genera, 20 species)
Series Bostrichiformia Forbes, 1926 (1 superfamily)
     Superfamily Bostrichoidea Latreille, 1802 (4 families)
          Family Dermestidae Latreille, 1804 (50 genera, 1200 species)
          Family Endecatomidae LeConte, 1861 (1 genus, 4 species)
          Family Bostrichidae Latreille, 1802 (90 genera, 570 species)
          Family Ptinidae Latreille, 1802 (230 genera, 2200 species)
Series Cucujiformia Lameere, 1938 (4 superfamilies)
     Superfamily Lymexyloidea Fleming, 1821 (1 family)
          Family Lymexylidae Fleming, 1821 (10 genera, 70 species)
     Superfamily Cleroidea Latreille, 1802 (12 families)
          Family Phloiophilidae Kiesenwetter, 1863 (1 genus, 1 species)
          Family Trogossitidae Latreille, 1802 (50 genera, 600 species)
```

See Löbl & Smetana (2007).

```
Family Chaetosomatidae Crowson, 1952 (4 genera, 12 species)
```

Family Metaxinidae Kolibáč, 2004 (1 genus, 1 species)

Family Thanerocleridae Chapin, 1924 (7 genera, 30 species)

Family Cleridae Latreille, 1802 (200 genera, 3400 species)

Family Acanthocnemidae Crowson, 1964 (1 genus, 1 species)

Family Phycosecidae Crowson, 1952 (1 genus, 4 species)

Family **Prionoceridae** Lacordaire, 1857 (3 genera, 160 species)

Family Mauroniscidae Majer, 1995 (5 genera, 26 species)

Family Melyridae Leach, 1815 (300 genera, 6000 species)

†Family Parandrexidae Kirejtshuk, 1994 (3 genera, 5 species)

Superfamily Cucujoidea Latreille, 1802 (37 families)

Family Boganiidae Sen Gupta and Crowson, 1966 (5 genera, 11 species)

Family Byturidae Gistel, 1848 (7 genera, 24 species)

Family Helotidae Chapuis, 1876 (5 genera, 107 species)

Family Protocucujidae Crowson, 1954 (1 genus, 7 species)

Family Sphindidae Jacquelin du Val, 1860 (9 genera, 59 species)

Family Biphyllidae LeConte, 1861(7 genera, 200 species)

Family Erotylidae Latreille, 1802 (260 genera, 3500 species)

Family Monotomidae Laporte, 1840 (33 genera, 250 species)

Family Hobartiidae Sen Gupta and Crowson, 1966 (2 genera, 6 species)

Family Cryptophagidae Kirby, 1826 (60 genera, 600 species)

Family Agapythidae Sen Gupta and Crowson, 1969 (1 genus, 1 species)

Family Priasilphidae Crowson, 1973 (3 genera, 11 species)

Family Phloeostichidae Reitter, 1911 (5 genera, 14 species)

Family Silvanidae Kirby, 1837 (58 genera, 500 species)

Family Cucujidae Latreille, 1802 (4 genera, 44 species)

Family Myraboliidae Lawrence and Britton, 1991 (1 genus, 13 species)

Family Cavognathidae Sen Gupta and Crowson, 1966 (1 genus, 9 species)

Family Lamingtoniidae Sen Gupta and Crowson, 1969 (1 genus, 3 species)

Family **Passandridae** Blanchard, 1845 (9 genera, 109 species)

Family **Phalacridae** Leach, 1815 (51 genera, 640 species)

Family **Propalticidae** Crowson, 1952 (2 genera, 30 species)

Family Laemophloeidae Ganglbauer, 1899 (37 genera, 430 species)

Family Tasmosalpingidae Lawrence and Britton, 1991 (1 genus, 2 species)

Family Cyclaxyridae Gimmel et al., 2009 (1 genus, 2 species)

Family Kateretidae Kirby, 1837 (14 genera, 95 species)

Family Nitidulidae Latreille, 1802 (350 genera, 4500 species)

Family Smicripidae G. H. Horn, 1880 (1 genus, 6 species)

Family Bothrideridae Erichson, 1845 (38 genera, 400 species)

Family **Cerylonidae** Billberg, 1820 (52 genera, 450 species)

Family Alexiidae Imhoff, 1856 (1 genus, 50 species)

Family Discolomatidae G. H. Horn, 1878 (16 genera, 400 species)

Family Endomychidae Leach, 1815 (130 genera, 1800 species)

Family Coccinellidae Latreille, 1807 (360 genera, 6000 species)

Family Corylophidae LeConte, 1852 (30 genera, 200 species)

Family Latridiidae Erichson, 1842 (28 genera, 1000 species)

Family Akalyptoischiidae Lord et. al, 2010 (1 genus, 24 species)<sup>6</sup>

†Family Sinisilvanidae Hong, 2002 (1 genus, 1 species)

Superfamily Tenebrionoidea Latreille, 1802 (28 families)

Family Mycetophagidae Leach, 1815 (18 genera, 130 species)

Family Archeocrypticidae Kaszab, 1964 (10 genera, 60 species)

Family Pterogeniidae Crowson, 1953 (7 genera, 26 species)

-

<sup>6.</sup> See Lord et al. (2010).

```
Family Ciidae Leach, 1819 (42 genera, 650 species)
```

Family Melandryidae Leach, 1815 (60 genera, 420 species)

Family Mordellidae Latreille, 1802 (100 genera, 1500 species)

Family Ripiphoridae Gemminger, 1870 (1853) (38 genera, 400 species)

Family **Zopheridae** Solier, 1834 (190 genera, 1700 species)

Family Ulodidae Pascoe, 1869 (14 genera, 30 species)

Family **Promecheilidae** Lacordaire, 1859 (7 genera, 20 species)

Family Chalcodryidae Watt, 1974 (3 genera, 15 species)

Family **Trachelostenidae** Lacordaire, 1859 (1 genus, 2 species)

Family Tenebrionidae Latreille, 1802 (2300 genera, 20000 species)

Family Prostomidae C. G. Thomson, 1859 (2 genera, 30 species)

Family Synchroidae Lacordaire, 1859 (3 genera, 8 species)

Family Stenotrachelidae C. G. Thomson, 1859 (7 genera, 19 species)

Family **Oedemeridae** Latreille, 1810 (100 genera, 500 species)

Family Meloidae Gyllenhal, 1810 (120 genera, 3000 species)

Family Mycteridae Oken, 1843 (29 genera, 160 species)

Family Boridae C. G. Thomson, 1859 (3 genera, 4 species)

Family Trictenotomidae Blanchard, 1845 (2 genera, 13 species)

Family **Pythidae** Solier, 1834 (7 genera, 23 species)

Family Pyrochroidae Latreille, 1807 (30 genera, 167 species)

Family Salpingidae Leach, 1815 (45 genera, 300 species)

Family Anthicidae Latreille, 1819 (100 genera, 3000 species)

Family Aderidae Winkler, 1927 (50 genera, 900 species)

Family Scraptiidae Gistel, 1848 (35 genera, 500 species)

Superfamily Chrysomeloidea Latreille, 1802 (7 families)

Family Vesperidae Mulsant, 1839 (17 genera, 75 species)

Family **Oxypeltidae** Lacordaire, 1868 (2 genera, 3 species)<sup>7</sup>

Family **Disteniidae** J. Thomson, 1861 (32 genera, 336 species)

Family Cerambycidae Latreille, 1802 (5232 genera, 30079 species)8

Family Megalopodidae Latreille, 1802 (30 genera, 350 species)<sup>9</sup>

Family **Orsodacnidae** C. G. Thomson, 1859 (3 genera, 40 species)<sup>9</sup>

Family Chrysomelidae Latreille, 1802 (2114 genera, 32500 species)

Superfamily Curculionoidea Latreille, 1802 (8 families)<sup>10</sup>

Family Nemonychidae Bedel, 1882 (20 genera, 70 species)

Family Anthribidae Billberg, 1820 (372 genera, 3900 species)

Family Belidae Schoenherr, 1826 (38 genera, 375 species)

Family Attelabidae (150 genera, 2500 species)

Family Caridae Thompson, 1992 (4 genera, 6 species)

Family **Brentidae** Billberg, 1820 (400 genera, 4000 species)

Family Curculionidae Latreille, 1802 (4600 genera, 51000 species)

†Family Ulyanidae Zherikhin, 1993 (2 genera, 3 species)

### Acknowledgements

We thank Petr Svacha (Vesperidae), Miguel Monne (Cerambycidae), Den Heffern (Cerambycidae), Antonio Santos-Silva (Disteniidae), and Chris Reid (Chrysomelidae) for assistance with the numbers of chrysomeloid families. RABL

Zootaxa 3148 © 2011 Magnolia Press · 207

Family **Tetratomidae** Billberg, 1820 (13 genera, 150 species)

<sup>7.</sup> See Kuschel (1955).

<sup>8.</sup> Number of genera was obtained by tripling the number of neotropical genera (1744) supplied by M. Monne (pers. com). Species number was obtained from Anonymous (2010).

<sup>9.</sup> See Hallan (2010)

<sup>10.</sup> See Oberprieler et al. (2007).

was supported in part by Foundation for Research, Science and Technology through the Defining New Zealand's Land Biota OBI.

#### Literature Cited

- Anonymous. (2010) Biolib.cz. http://www.biolib.cz/en/taxon/id10991/ [accessed 6 January 2011].
- Beutel, R. G., & Leschen, R.A.B. (2005) Handbook of Zoology, Volume IV Arthropoda: Insecta, Part 38, Evolution and Systematics Coleoptera (Archostemata, Adephaga, Myxophaga, Polyphaga part.). Walter de Gruyter, Berlin. 567 p.
- Bouchard, B, Bousquet, Y, Davies, A.E., Alonso-Zarazaga, M.A., Lawrence, J.F., Lyal, C.H.C., Newton, A.F., Reid, C.A.M., Schmitt, M., Ślipiński, S.A., Smith, A.B.T. (2011) Family-group names in Coleoptera. *Zookeys*, 88, 1–972.
- Fikáček, M., Šípková H. (2009) New Asian *Hydroscapha*, with comments on male-female association of co-occuring species (Coleoptera, Myxophaga, Hydroscaphidae) *Zootaxa*, 2286, 31–48.
- Ge, S.-Q., Friedrich, F. & Beutel, R. G. (2010) On the systematic position and taxonomic rank of the extinct myxophagan †*Haploche-lus* (Coleoptera). *Insect Systematics & Evolution*, 41, 329–338.
- Hallan, J. (2010) Synopsis of the described Coleoptera of the world. http://insects.tamu.edu/research/collection/hallan/test/Arthropoda/Insects/Coleoptera/Family/Coleoptera1.htm [accessed 6 January 2011].
- Kirejtshuk, A.G., Ponomarenko, F.G. & Zherikhin, V.V. (2010) *Catalogue of fossil Coleoptera*. http://www.zin.ru/animalia/coleoptera/eng/paleosys.htm. [Accessed November 2010].
- Kuschel, G. (1955) Una nueva espécie de Cheloderus Castelnau. Revista Chilena de Entomología, 4, 251-254.
- Lawrence, J.F. & Newton, A. F. (1995) Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). *In:* Pakaluk, J. & Ślipiński, S. A. (Eds), *Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson.* Muzeum Instytut Zoologii PAN, Warszawa, pp. 779–1006.
- Leschen, R.A.B., Beutel, R.G., & Lawrence, J.F. (2010) Handbook of Zoology, Coleoptera Volume 2: Morphology and Systematics (Elateroidea, Bostrichformia, Cucujiformia partim). Walter de Gruyter, Berlin. 786 p.
- Löbl, I. & Smetana A. (2007) Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea and Cucujoidea. Apollo Books, Stenstrup. 935 p.
- Lord, N.P., Hartley, C.S., Lawrence, J.F., McHugh, J.V., Whiting, M.F., & Miller, K.B. (2010) Phylogenetic analysis of the Minute Brown Scavenger Beetles (Coleoptera: Latridiidae), and recognition of a new beetle family, Akalyptoischionidae, fam. n. (Coleoptera: Cucujoidea). *Systematic Entomology*, 35, 753–763.
- Oberprieler, R.G., Marvaldi, A.E. & Anderson, R.S. (2007) Weevils, weevils, weevils everywhere. *In:* Zhang, Z.-Q. & Shear, W.A. (Eds.) *Linnaeus Tercentenary: Progress in Invertebrate Taxonomy. Zootaxa*, 1668, 491–520.

## Author names and contact addresses

- S.A. Slipinski, CSIRO Ecosystem Sciences, GPO Box 1700, Canberra, ACT, 2601, Australia
- R.A.B. Leschen, New Zealand Arthropod Collection, Private Bag 92170, Auckland, New Zealand (corresponding author; leschenr@landcareresearch.co.nz)
- J.F. Lawrence, CSIRO Ecosystem Sciences, GPO Box 1700, Canberra, ACT, 2601, Australia